

Aberystwyth University

Polyphenol oxidase in leaves

Boeckx, Tinne; Winters, Ana; Webb, Judith; Kingston-Smith, Alison

Published in:

Journal of Experimental Botany

DOI:

[10.1093/jxb/erv141](https://doi.org/10.1093/jxb/erv141)

Publication date:

2015

Citation for published version (APA):

Boeckx, T., Winters, A., Webb, J., & Kingston-Smith, A. (2015). Polyphenol oxidase in leaves: Is there any significance to the chloroplastic localization? *Journal of Experimental Botany*, 66(12), 3571-3579.
<https://doi.org/10.1093/jxb/erv141>

Document License

CC BY

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400

email: is@aber.ac.uk



REVIEW PAPER

Polyphenol oxidase in leaves: is there any significance to the chloroplastic localization?

Tinne Boeckx^{*}, Ana L. Winters, K. Judith Webb and Alison H. Kingston-Smith[†]

Institute of Biological, Environmental and Rural Sciences (IBERS), Penglais Campus, Aberystwyth University, Aberystwyth SY23 3FG, UK

^{*} Present address: Division of Plant and Crop Sciences, University of Nottingham, Sutton Bonington, Nottingham, UK

[†] To whom correspondence should be addressed. E-mail: ahk@aber.ac.uk

Received 19 December 2014; Revised 27 February 2015; Accepted 2 March 2015

Abstract

Polyphenol oxidase (PPO) catalyses the oxidation of monophenols and/or *o*-diphenols to *o*-quinones with the concomitant reduction of oxygen to water which results in protein complexing and the formation of brown melanin pigments. The most frequently suggested role for PPO in plants has been in defence against herbivores and pathogens, based on the physical separation of the chloroplast-localized enzyme from the vacuole-localized substrates. The *o*-quinone–protein complexes, formed as a consequence of cell damage, may reduce the nutritional value of the tissue and thereby reduce predation but can also participate in the formation of structural barriers against invading pathogens. However, since a sufficient level of compartmentation-based regulation could be accomplished if PPO was targeted to the cytosol, the benefit derived by some plant species in having PPO present in the chloroplast lumen remains an intriguing question. So is there more to the chloroplastic location of PPO? An interaction between PPO activity and photosynthesis has been proposed on more than one occasion but, to date, evidence either for or against direct involvement has been equivocal, and the lack of identified chloroplastic substrates remains an issue. Similarly, PPO has been suggested to have both pro- and anti-oxidant functions. Nevertheless, several independent lines of evidence suggest that PPO responds to environmental conditions and could be involved in the response of plants to abiotic stress. This review highlights our current understanding of the *in vivo* functions of PPO and considers the potential opportunities it presents for exploitation to increase stress tolerance in food crops.

Key words: Abiotic stress, photosynthesis, polyphenol oxidase, secondary metabolism.

Introduction

Over the past five decades, yields of wheat and maize crops have decreased by an estimated 1–2% per decade, affecting food supplies for both humans and livestock (IPCC, 2014). In order to feed a growing human population, global food production must be maintained or, preferably, increased, under increasingly unstable climatic conditions and rising temperatures (IPCC, 2014). New approaches are needed to meet this major agricultural challenge whilst using existing or even reduced tracts of agricultural land (Foresight, 2011).

A key agricultural target is to improve the yields of crops growing under periods of abiotic, as well as biotic, stress. The enzyme polyphenol oxidase (PPO) is found in most plant species and the foliar expressed gene products may have a role in either acclimation or short-term response to stress, indicated by circumstantial evidence such as enzyme localization and its response to environmental factors. PPO has been reported in all land plants surveyed to date with the exception of *Arabidopsis*. By contrast, no PPO-like sequences have been reported in chlorophytes (green algae; Tran *et al.*, 2012).

This study showed that the size of PPO gene families varied widely with numbers of PPO genes ranging from one to 13 in the 18 genomes analysed. It is postulated that the occurrence of this enzyme is strongly correlated with the emergence of land plants suggesting a role in adaption to abiotic stress associated with a dry/non-aquatic environment. However, as yet, there is no conclusive evidence for the existence of an underlying mechanism explaining the relationship between PPO and abiotic stress; indeed it is unclear if the presence of PPO activity is beneficial or detrimental to the plant (Mayer, 2006). While PPO activity can be related to the accumulation of reactive oxygen species (ROS; Thipyapong et al., 2004b; Mayer, 2006) and overall redox potential values (Webb et al., 2014), its presence could also be beneficial as a proposed oxygen buffer (Vaughn and Duke, 1984) or by down-regulating photosynthesis (Trebst and Depka, 1995).

The current state of what is often contradictory hypothetical and experimental evidence regarding the potential of PPO in leaves to confer an advantage in crop production, particularly during periods of abiotic stress, such as drought, heat, and cold is examined here.

Fundamentals of PPO biochemistry

PPO enzymes from plants comprise three domains including an N-terminal plastid transit peptide, a highly conserved type-three copper centre, and a C-terminal region (Tran et al., 2012). The family of PPO enzymes catalyse the oxidation of monophenols and/or *o*-diphenols to *o*-quinones. PPOs are widely distributed in bacteria, animals, plants, and fungi (Mayer, 2006; Tran et al., 2012) but are often confused with another subclass of phenol oxidases, the

laccases [benzenediol: oxygen oxidoreductase [EC 1.10.3.2] or *p*-diphenol oxidase], which oxidize a broad range of *o*-, *m*-, and *p*-diphenols (Griffith, 1994). Plant laccases are mostly extracellular proteins with 22–45% glycosylation (Solomon et al., 1996) whereas PPOs are intracellular proteins and the degree of glycosylation is unclear (Steffens et al., 1994). Both are multicopper oxidases, however, PPOs have a coupled binuclear type-3 copper centre and laccases have a trinuclear cluster of four copper ions (Fig. 1; Solomon et al., 1996).

Subclasses of PPO enzymes are tyrosinases or catecholases according to the absence or presence of cresolase or monophenolase activity (Fig. 2.). Tyrosinases first catalyse the oxidation of monophenols to *o*-diphenols (monophenolase activity; EC 1.14.18.1), and subsequently catalyse the *o*-diphenol to *o*-quinone reaction (catecholase activity; EC 1.10.3.1). Catecholases on the other hand are *o*-diphenol specific and so are only able to catalyse the oxidation of *o*-diphenols to *o*-quinones (Steffens et al., 1994). Both subclasses are, however, commonly referred to as PPO (Mayer and Harel, 1979; Marusek et al., 2006) since the poor characterization of these enzymes could mean that some catecholases are, in fact, tyrosinases in which the monophenolase activity has not yet been observed (Solomon et al., 1996), as detection of monophenolase activity requires highly specific conditions compared with catecholase activity (Yoruk and Marshall, 2003).

The primary reaction products of PPO activity are the *o*-quinones, which are highly reactive and will covalently modify and cross-link proteins to form brown melanin pigments (Steffens et al., 1994; Kroll et al., 2000; Kroll and Rawel, 2001). An example of this is the browning of apples and potatoes, seen shortly after wounding or cutting.




Fig. 1. Active sites of (A) tyrosinase/ catecholase and (B) laccase enzymes. Both (A) and (B) contain a binuclear type 3 copper centre and (B) also includes mononuclear type 1 and 2 copper centres.




Fig. 2. Schematic illustration of the mechanism of PPO including the structures of the *o*-diphenol and monophenol substrates for the catecholase and monophenolase reactions.

PPO-mediated generation of *o*-quinones has also been implicated in the indirect generation of ROS as secondary reaction products (Steffens *et al.*, 1994). Although the precise mechanism remains to be established, reverse disproportionation of *o*-quinones can result in the formation of cytoplasmic semiquinone radicals (O'Brien, 1991; Thipyapong *et al.*, 1997). Interaction between these radicals and O₂ will result in the generation of superoxide anions and the regeneration of *o*-quinone (O'Brien, 1991). Superoxide anions are very unstable and will quickly dismutate, either enzymatically via superoxide dismutase, or non-enzymatically, to form hydrogen peroxide (Grant and Loake, 2000). A Fenton reaction between divalent metal ions such as iron (II) and the relatively stable hydrogen peroxide can result in the generation of extremely reactive hydroxyl radicals. Accumulation of these cytotoxic ROS needs to be under tight control as oxidative modifications including protein cross-linking, lipid peroxidation, and damage to nucleic acids may ultimately inflict cell death (Grant and Loake, 2000; Bhattacharjee, 2005; Gill and Tuteja, 2010; Foyer and Noctor, 2012). However, theoretically, PPO could also contribute to decreasing the amount of oxygen locally available through the reduction of O₂ to water (Yoruk and Marshall, 2003).

Regulation of PPO is complex and the enzyme can be present in both an active and a latent (an inactive, often

precursor form) state in the same source material (Mayer and Harel, 1979). Following transport to the lumen and the cleavage of the N-terminal transit peptide, PPO is initially present as a two-domain protein consisting of a copper binding site and a C-terminal domain (Flurkey and Inlow, 2008). A more detailed discussion, including consideration of the 3-D structure of catechol oxidases, can be found in Gerdemann *et al.* (2002). The C-terminal is linked via a highly flexible random peptide structure which is proposed to cover the active site and undergo conformational change under certain conditions (Leufken *et al.*, 2014). Flurkey and Inlow (2008) have reviewed evidence for C-terminal proteolytic processing of latent PPO to the active form which has been demonstrated for *Vicia faba*, *Vitis vinifera*, and *Ipomoea batatas* PPOs. The degree of latency is not universal and differs with plant species as well as tissue type. For instance, PPO activity was detected in both the active and latent forms in root tissues of red clover (*Trifolium pratense*), but only in the latent form in white clover (*Trifolium repens*); this contrasts with aerial tissues where PPO activity was detected in both active and latent forms in both red and white clover (Webb *et al.*, 2013). Protein in this latent state is not only activated by proteolytic cleavage but also by chemically inducing conformational changes to the latent enzyme. *In vitro*, treatments that are effective in activating latent PPO include exposure to fatty acids, proteolysis

(trypsin), mild heat, acid and base shocks, and detergents such as SDS and ammonium sulphate (Tolbert, 1973; Steffens *et al.*, 1994; Jiménez and García-Carmona, 1996; Yoruk and Marshall, 2003). Leufken *et al.* (2014) observed that the C-terminal domain determines the pH optimum of plant PPOs in non-proteolytic activated enzyme and they postulate that non-proteolytic activation also occurs *in planta*. *In vivo*, activation of the latent PPO pool could occur as a result of direct interaction between the enzyme and its substrates. Winters *et al.* (2008) have demonstrated the potential to activate latent PPO from red clover in the presence of its endogenous *o*-diphenols substrates. It has been proposed that *o*-diphenol-mediated activation is an indirect mechanism of activation, with the resulting *o*-quinones interacting with the latent PPO pool, thereby altering their structure and exposing the active sites (Winters *et al.*, 2008). This is presumably the mechanism occurring upon tissue damage, either as a result of herbivory (Lee *et al.*, 2009), or of senescence-associated cell disruption, as reported by Meyer and Biehl (1980) who observed an increase in phenolase activity and a concomitant decrease of the latent form of PPO during leaf ageing in spinach (*Spinacia oleracea*). More recently, Molitor *et al.* (2013) identified a putative quinone binding site in the PPO enzyme, aurone synthase, from *Coreopsis grandiflora*, which they propose is responsible for the observed allosteric activation of latent PPO.

The conundrum of PPO compartmentation

Arnon (1949) provided some of the earliest evidence for the intracellular location of PPO in spinach beet (*Beta vulgaris*) chloroplasts. This has been followed up by detailed investigations revealing that, in leaves, PPO is specifically located in the lumen or loosely attached to the luminal side of the thylakoid membrane (Tolbert, 1973; Mayer and Harel, 1979; Sommer *et al.*, 1994) in the vicinity of photosystems I and II (Lax and Vaughn, 1991). The mechanism by which this is achieved was revealed by Sommer *et al.* (1994). They demonstrated that the nuclear-encoded sequences were likely to behave as characteristic thylakoid-targeted proteins which use the light-generated thylakoid pH gradient as the energy source to transport the intermediates of stromal proteins across the thylakoid membrane and into the lumen, otherwise known as the Δ pH pathway (Keegstra and Cline, 1999). More recently, PPOs have also been identified which lack the chloroplast targeting sequence (Tran *et al.*, 2012) and have been found in the cytosol (Nakayama *et al.*, 2000; 2001) and in the vacuole (Ono *et al.*, 2006).

In contrast to the mostly chloroplastic location of the PPO protein, phenolic compounds are generally confined to the vacuoles (Mayer and Harel, 1979; Vaughn and Duke, 1984). This includes those compounds recognized as substrates for PPO. To date, potential PPO substrates have been identified within the anthocyanin, flavanol, flavone, flavonol, and isoflavonoid subclasses of flavonoid polyphenols and hydroxybenzoic acid and hydroxycinnamic acid subclasses of phenolic acids based on enzyme assays or structural comparison to

confirmed substrates (Parveen *et al.*, 2010). Given the physical separation of PPO enzymes from their substrates, it is commonly accepted that the PPO enzyme–substrate interaction requires the destruction of cell compartmentation, as a result of wounding for example. It is therefore not surprising that PPO enzyme activity has usually been related to arthropod (Kowalski *et al.*, 1992) or pathogen defence mechanisms (Li and Steffens, 2002; Thipyapong *et al.*, 2004a) as *o*-quinone protein complexes can decrease the nutritional value of the tissue (Felton *et al.*, 1989, Thipyapong *et al.*, 2004a) and/or ROS (secondary PPO reaction products) could trigger defence pathways (Kowalski *et al.*, 1992; Thipyapong, 2007).

At face value, this physical separation of enzyme and substrate appears logical, although somewhat wasteful energetically to target mature PPO protein to the thylakoid lumen while sufficient compartmentation away from vacuolar substrates could also be achieved by targeting PPO to the cytosol. Therefore, considering the widespread occurrence of this trait in higher plants, even in the absence of detectable substrate (e.g. *Medicago sativa*; Sullivan *et al.*, 2008), this highly specific localization of PPO could indicate that it confers a distinct advantage. However, a chloroplastic role for PPO is far from clear. For PPO activity to have a function in undamaged tissues it is necessary for the enzyme to have ready access to a suitable substrate in the chloroplast. Typical PPO substrates are *o*-diphenols because of their readily oxidizable OH-groups (Martinez and Whitaker, 1995; Parveen *et al.*, 2010). Most recognized PPO substrates are appointed to just two classes of polyphenols; the phenolic acids and the flavonoids (Parveen *et al.*, 2010). Although the presence of polyphenols and flavonoids have been reported in chloroplasts (Satô, 1966; Halliwell, 1975; Saunders and McClure, 1976a, b; Agati *et al.*, 2007; Liu *et al.*, 2009), to our knowledge, only catechin has so far been reported as a substrate for PPO in the mesophyll chloroplasts of tea (Subramanian *et al.*, 1999; Liu *et al.*, 2009). The identification of further potential monophenolic and/or *o*-diphenolic PPO substrates in chloroplasts will be paramount in order to demonstrate an *in vivo* function of PPO in undamaged tissue.

The relationship between PPO and environment

As well as evidence in favour of the involvement of PPO in plant defence against biotic stressors, several independent lines of evidence implicate the chloroplastic location of PPO in an, as yet uncharacterized, contribution to the response of plants to abiotic stress, potentially mediated by altering the cellular balance of ROS production/degradation. It has been suggested that an acclimation mechanism exists whereby the oxidation of accumulated phenolics is inhibited when plants are subjected to extreme temperatures (Rivero *et al.*, 2001) or drought (Sofa *et al.*, 2005; Lee *et al.*, 2007). Indeed, experimentally-imposed conditions of cold, heat, and drought have been shown to result in a significant increase in total phenolic compounds compared with the controls (Rivero *et al.*, 2001; Sofa *et al.*, 2005; Lee

et al., 2007). Concomitantly, oxidation of these accumulated phenolics was proposed to be inhibited by significant decreases in PPO (Rivero et al., 2001; Sofo et al., 2005) and peroxidase activities (Rivero et al., 2001) and through the significant activation of enzymatic ROS scavengers such as ascorbate peroxidase (Sofo et al., 2005; Lee et al., 2007) and superoxide dismutase (Sofo et al., 2005) compared with the controls. Hence the suggestion that a decrease in PPO activity following abiotic stress was associated with improved antioxidant capacity (Sofo et al., 2005). This was supported by the work of Thipyapong et al. (2004b) which showed that suppression of PPO increased the drought tolerance of tomato. However, a conflicting result was obtained when a drought treatment on white clover (*T. repens*) significantly increased PPO activity after 7 d (Lee et al., 2007). Interestingly, PPO is implicated in the adaption of resurrection plants to desiccation and rehydration; Veljovic-Jovanovic et al. (2008) observed an increase in PPO of several fold when *Ramonda serbica* leaves were subjected to near-complete water loss.

Given the contrasting responses of PPO activity to environmental conditions, it is not surprising that the potential impact of altered PPO activity on plant development, phenotype, and yield is currently unclear. In tomato (*Solanum esculentum*), the alteration of PPO activity by silencing did not affect plant development, total leaf area or shoot and root dry weights under optimal growth conditions when compared with non-transformed controls (Thipyapong et al., 2004b). Similarly, transgenic RNAi lines and wild-types of red clover (*T. pratense*) did not differ significantly in growth and leaf nitrogen content under optimal growth conditions (Webb et al., 2013). Alternatively, a clear effect of PPO silencing was observed in walnut plants (*Juglans regia*) which developed spontaneous necrotic lesions in the leaves even when not challenged by pathogens (Araji et al., 2014), suggesting increased susceptibility to oxidative stress. Furthermore, while no obvious phenotypic differences between wild-type red clover (*T. pratense* cv. Milvus) and a low PPO mutant were reported (Winters et al., 2008), a field study in Aberystwyth (UK) reported a higher dry matter yield from fields seeded with wild-type red clover (5.78 tonnes DM ha⁻¹) than when seeded with the low PPO mutant (5.40 tonnes DM ha⁻¹) (R Fychan, unpublished data; 0.4 ha were sown with each of red clover cv. Milvus and the PPO mutant in 2009 and the total dry matter weight of the above-ground matter harvested in May 2010 was determined). Notably, both growth conditions and developmental stage influences the import of PPO into the chloroplasts (Sommer et al., 1994) with corresponding effects on PPO activity (Mayer and Harel, 1979; Webb et al., 2013), possibly accounting for the observed changes in PPO activity in red clover during a growing season (Fig. 3; Fothergill and Rees, 2006). The latter demonstrates the impact of seasonal variation, with a peak in PPO activity in the winter months, during which the combination of high light and the relatively low demand for fixed carbon results in a high risk of photoinhibition and the associated oxidative stress.




Fig. 3. Seasonal total PPO activity for July 2004–June 2005. Mean of six red clover (*Trifolium pratense*) cultivars are shown \pm SE (original data courtesy of M Fothergill; Fothergill and Rees, 2006).

Could PPO be involved in photosynthesis?

A potential role for PPO in photosynthesis has been speculated on previously (Mayer and Harel, 1979; Vaughn and Duke, 1984). Although theoretically plausible, evidence supporting or countering this hypothesis is still poor. Observations in support of the involvement of PPO in photosynthesis include: (i) the correlation between PPO activity and chloroplasts evolving high levels of O₂ (Vaughn and Duke, 1984), (ii) the association of PPO protein with the photosystems (Lax and Vaughn, 1991; Sheptovitsky and Brudvig, 1996), (iii) the inhibition of cyclic and/or non-cyclic photophosphorylation by phenolic compounds (Neumann and Drechsler, 1967), the implication being that PPO activity could prevent such inhibition by oxidation of these potential substrates, (iv) the independence of increases and decreases of substrate level and catecholase activity during growth and development (Ben-Shalom et al., 1977; Winters et al., 2008; Webb et al., 2014), and, finally (v) the modulation of PPO activity by environmental effects such as extremes of temperature, drought, and time of year (Rivero et al., 2001; Thipyapong et al., 2004b; Sofo et al., 2005; Fothergill and Rees, 2006; Lee et al., 2007).

There have been several suggestions as to a mechanism by which PPO could directly influence photosynthesis, including it functioning as an oxygen buffer (Mayer and Harel, 1979; Vaughn and Duke, 1984) or interacting with the Mehler–peroxidase, or water–water, cycle (Tolbert, 1973) to facilitate reactive oxygen scavenging (Fig. 4). The possibility that PPO modulates available oxygen is plausible given the requirement for O₂ during PPO-catalysed oxidation of phenolic compounds to *o*-quinones and H₂O (Steffens et al., 1994), while the close association with the photosystems could provide a source of sufficient reducing power to regenerate *o*-diphenol by reducing *o*-quinones (Halliwell, 1975; Vaughn and Duke, 1984). The limitations to this hypothesis are the lack of definitively identified chloroplast substrates and the relative slow rate of this catalysis compared with the speed of photosynthesis. Interaction with the Mehler–peroxidase cycle was




Fig. 4. The potential interaction of PPO with photosynthesis (i) by acting as an oxygen buffer in the lumen to prevent O_2^- formation, and (ii) to buffer NADPH accumulation in the stroma to prevent over-reduction of the photosystems. GSH (GSSG), reduced (oxidized) glutathione; DHA, dehydroascorbate; GR, glutathione reductase; DHAR, dehydroascorbate reductase.

proposed as quinones can also serve as hydrogen acceptors, with the reduced quinones being re-oxidized by PPO. In this way, a pseudo-cyclic electron transport would occur without a net oxygen change (Trebst et al., 1963; Tolbert, 1973), which is how the water–water cycle is believed to operate (Asada, 1999). There is, however, no concrete evidence to support either of these hypotheses and the water–water cycle and ROS scavenging are already well described processes (Asada, 1999; Apel and Hirt, 2004). Indeed, contrary to the above hypotheses, a study of tomato plants in which PPO was suppressed by transformation showed that the transformants actually performed better under conditions designed to impose photoinhibition than did the untransformed plants (Thipyapong et al., 2004b).

A related mechanism could potentially counter over-reduction due to the accumulation of high NAD(P)H/NAD(P) ratios which can lead to the inactivation of photosynthetic electron transport (Foyer et al., 2012). An *o*-dihydroxyphenol–ascorbate reduction/oxidation cycle linked with the ascorbate–glutathione cycle, which involves oxidation of NAD(P)H, could provide a mechanism for preventing over-reduction under conditions of decreased CO_2 fixation. This could potentially be catalysed by low pH-activated latent PPO (see below) in high-light conditions with elevated levels of lumen O_2 . While this reduces the requirement for cyclic electron transport, and quinones react readily with ascorbic acid, the extent to which ascorbate would participate in this, compared with other luminal processes, is unclear (Tóth et al., 2013).

A proposed alternative explanation is that PPO is more important for the dark reactions in the thylakoid lumen than those in the light, i.e. when O_2 is low (Sheptovitsky and Brudvig, 1996). This was suggested since an acidic environment is created in the light in which the PPO enzyme, with pH optima of 8, would not be expected to be active (Sheptovitsky and Brudvig, 1996). However, this simple theory seems doubtful as various previous and subsequent studies reported pH optima of between 4 and 8 for PPO (Tolbert, 1973; Rocha and Morais, 2001; Yoruk and Marshall, 2003). This variability may be explained by the work of Leufken et al. (2014) who demonstrated that conformational change in the C-terminus can determine pH optima. Furthermore, latent PPO is known

to be activated by low pH (Steffens et al., 1994; Winters et al., 2003; Schmitz et al., 2008), which correlates well with the decrease in lumen pH following the illumination of chloroplasts. Plus, it would appear that a source of reducing power is necessary for intra-chloroplastic PPO activity; the oxidation of *p*-coumaric acid to caffeic acid observed in illuminated isolated chloroplasts did not occur in the dark unless ascorbate or NADPH was added (Halliwell, 1975). These observations add further support for the role of PPO in high-light conditions involving ascorbate and/or NAD(P)H.

In many respects however, a direct role for PPO in the regulation of photosynthesis may be unrealistic given the flux through the respective pathways (Vinyard et al., 2013). It is therefore more likely that PPO has an indirect role in photosynthesis. As PPO monophenolase activity could theoretically catalyse the conversion of *p*-coumaric acid to caffeic acids (Vaughn and Duke, 1984), which is one of the initial steps in the phenylpropanoid pathway of phenolic compound biosynthesis, a role via secondary metabolism is possible. This is supported by recently published work in which the silencing of PPO gene expression in walnut plants altered the metabolite profile of the leaves, notably those involved in tryptophan and tyrosine metabolism (Araji et al., 2014). It was argued that the observed endogenous increases in tyramine and decreases in DOPA in PPO silenced plants occurred because, normally, PPO catalysed the *o*-hydroxylation of tyrosine to DOPA and tyramine to dopamine. So far, no enzymes have been characterized for these reactions in walnut. The observation that the endogenous application of tyramine to wild-type leaves caused the same necrotic phenotype seen in the PPO-silenced plants, supports a fundamental role for PPO in tyrosine metabolism in walnut. Also, the product of a PPO transcript from *Coreopsis grandiflora* has been demonstrated to be involved in aurone formation (Kaintz et al., 2014) and further supports a potential role for PPO in secondary metabolism. Several reports have theorized that the antioxidant capacity of phenolics (Pietta, 2000; Parveen et al., 2010) is indicative of their potential as radical scavengers (Neill and Gould, 2003; Agati et al., 2007) and even as photochemical energy dissipaters, the focus being on the phenolic acid subclass of hydroxycinnamic acids (chlorogenic acid specifically; Grace and Logan, 2000) and the anthocyanin subclass of the

flavonoids (Neill and Gould, 2003). Therefore, by regulating the availability of phenolics, PPO would indirectly affect the photoprotective capacity of photosynthetic cells independently of processes such as the Mehler–peroxidase cycle. However, this is not a uniform response and up-regulation of phenylpropanoid metabolism is not always accompanied by decreases in PPO activity (Rivero *et al.*, 2001; Sofo *et al.*, 2005; Fothergill and Rees, 2006; Lee *et al.*, 2007).

Future challenges

PPO is an enigmatic enzyme with many possibilities but few certainties. Confusingly, the data suggest that PPO activity can confer both a productive advantage and be associated with an increased risk of oxidative damage. While PPO activity can be associated with non-enzymatic ROS scavenging involving flavonoid and phenolic acid substrates (Apel and Hirt, 2004; Parveen *et al.*, 2010), a role for PPO in plant function may also be associated with its pro-antioxidant activity through the generation of secondary reaction products, including ROS (O'Brien, 1991; Thipyapong *et al.*, 1997), or may even involve localized effects in cellular differentiation/death, where concentrations may be locally high and critical but not easily measurable, for example, in nodules (Webb *et al.*, 2014) and walnut (Araji *et al.*, 2014). The evidence indicates that PPO performs different roles in different plant species and possibly multiple roles in plants with large PPO gene families. It is therefore important that fundamental questions, such as whether the *in vivo* role involves a pro- or anti-oxidant, would greatly increase our understanding of this enzyme and clarify future opportunities to exploit its function for increased and sustainable crop production. Consideration of possible relationships between PPO activity and photosynthesis are clearly relevant to the current issues of food security. The global population is increasing rapidly and is due to reach nine billion in 2050, creating a pressing need to optimize sustainable food production, with careful attention on crops for both human consumption and animal feed (Kingston-Smith *et al.*, 2013). To date, results of research into a possible role for PPO in photosynthesis has yielded equivocal results, possibly because the focus has been on looking for a direct effect on regulation or mitigation of photochemistry. It is proposed here that a re-examination of possible indirect effects of PPO on photosynthetic performance under abiotic stress is appropriate to current global challenges. Tools such as the identification of mutants (Lee *et al.*, 2004; Winters *et al.*, 2008), plant material genetically altered in PPO composition (Sullivan *et al.*, 2004; Thipyapong *et al.*, 2004b; Araji *et al.*, 2014; Webb *et al.*, 2014), and *in vitro* cloning to continue the identification of still unknown catalysts of phenolic compound biosynthesis (Sullivan and Zarnowski, 2010) plus an increased use of unbiased analytical techniques for metabolic analysis holds the promise of finding explanations to some of the unexplained aspects of PPO biochemistry. These analyses should determine the extent to which endogenous PPO activity has the potential to improve photosynthetic performance under abiotic stress conditions and indicate whether

increased activities of this enzyme should be included as a desirable leaf trait in plant breeding programmes designed to increase the yield of food crops.

Acknowledgements

This work was supported by an IBERS postgraduate studentship (TB) and a BBSRC Institute Strategic Programme Grant (BBS/E/W/10964A01; AK-S).

References

- Agati G, Matteini P, Goti A, Tattini M. 2007. Chloroplast-located flavonoids can scavenge singlet oxygen. *New Phytologist* **174**, 77–89.
- Apel K, Hirt H. 2004. Reactive oxygen species: metabolism, oxidative stress and signal transduction. *Annual Review of Plant Biology* **55**, 373–399.
- Araji S, Grammer TA, Gertzen R, *et al.* 2014. Novel roles for the polyphenol oxidase enzyme in secondary metabolism and the regulation of cell death in walnut (*Juglans regia*). *Plant Physiology* **164**, 1191–1203.
- Arnon DL. 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology* **24**, 1–15.
- Asada K. 1999. The water–water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons. *Annual Review of Plant Physiology and Plant Molecular Biology* **50**, 601–639.
- Ben-Shalom N, Kahn V, Harel E, Mayer AM. 1977. Olive catechol oxidase: changes during fruit development. *Journal of the Science of Food and Agriculture* **28**, 545–550.
- Bhattacharjee S. 2005. Reactive oxygen species and oxidative burst: roles in stress, senescence and signal transduction in plants. *Current Science* **89**, 1113–1121.
- Felton GW, Donato K, Del Vecchio RJ, Duffey SS. 1989. Activation of plant foliar oxidases by insect feeding reduces nutritive quality of foliage for noctuid herbivores. *Journal of Chemical Ecology* **15**, 2667–2694.
- Flurkey WH, Inlow JK. 2008. Proteolytic processing of polyphenol oxidase from plants and fungi. *Journal of Inorganic Biochemistry* **102**, 2160–2170.
- Foresight. 2011. *The future of food and farming: final project report*. London: The Government Office for Science.
- Fothergill M, Rees ME. 2006. Seasonal differences in polyphenol oxidase activity in red clover. In: Wachendorf M, Helgadottir A, Parente G, eds. *Sward dynamics, N-flows and forage utilisation in legume-based systems*. Proceedings of COST Workshop, Grado, Italy, 10–12 November 2005, 141–144.
- Foyer CH, Noctor G. 2012. Managing the cellular redox hub in photosynthetic organisms. *Plant, Cell and Environment* **35**, 199–201.
- Foyer CH, Neukermans J, Queval G, Noctor G, Harbinson J. 2012. Photosynthetic control of electron transport and the regulation of gene expression. *Journal of Experimental Biology* **63**, 1637–1661.
- Gerdemann C, Eicken C, Krebs B. 2002. The crystal structure of catechol oxidase: new insight into the function of type-3 copper proteins. *Accounts of Chemical Research* **35**, 183–191.
- Gill SS, Tuteja N. 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry* **48**, 909–930.
- Griffith GW. 1994. Phenoloxidases. In: Martinelli SD, Kinghorn JR, eds. *Progress in industrial microbiology, Vol. 29: Aspergillus nidulans: 50 years on*. Amsterdam: Elsevier Science Publishers, 763–788.
- Grace SC, Logan BA. 2000. Energy dissipation and radical scavenging by the plant phenylpropanoid pathway. *Philosophical Transactions of the Royal Society London, Series B* **355**, 1499–1510.
- Grant JJ, Loake GJ. 2000. Role of reactive oxygen intermediates and cognate redox signaling in disease resistance. *Plant Physiology* **124**, 21–29.
- Halliwell B. 1975. Hydroxylation of *p*-coumaric acid by illuminated chloroplasts: The role of superoxide. *European Journal of Biochemistry* **55**, 355–360.

- IPCC (Intergovernmental Panel on Climate Change).** 2007. Summary for policy makers. In: Field CB, Barros VR, Dokken DJ *et al.*, eds. *Climate change 2014: impacts, adaptation and vulnerability*. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, UK and New York: Cambridge University Press, 1–32.
- Jiménez M, García-Carmona F.** 1996. The effect of sodium dodecyl sulphate on polyphenol oxidase. *Phytochemistry* **42**, 1503–1509.
- Kaintz C, Molitor C, Thill J, Kampatsikas I, Michael C, Halbwirth H, Rempel A.** 2014. Cloning and functional expression in *E. coli* of a polyphenol oxidase transcript from *Coreopsis grandiflora* involved in aurone formation. *FEBS Letters* **588**, 3417–3426.
- Keegstra K, Cline K.** 1999. Protein import and routing systems of chloroplasts. *The Plant Cell* **11**, 557–570.
- Kingston-Smith AH, Marshall AH, Moorby JM.** 2013. Breeding for genetic improvement of forage plants in relation to increasing animal production with reduced environmental footprint. *Animal* **7**, Supplement 1, 79–88.
- Kowalski SP, Eannetta NT, Hirzel AT, Steffens JC.** 1992. Purification and characterization of polyphenol oxidase from glandular trichomes of *Solanum berthaultii*. *Plant Physiology* **100**, 677–684.
- Kroll J, Rawel HM.** 2001. Reactions of plant phenols with myoglobin: influence of chemical structure of the phenolic compounds. *The Journal of Food Science* **66**, 48–58.
- Kroll J, Rawel HM, Seidelmann N.** 2000. Physicochemical properties and susceptibility to proteolytic digestion of myoglobin-phenol derivatives. *Journal of Agricultural and Food Chemistry* **48**, 1580–1587.
- Lax AR, Vaughn KC.** 1991. Colocalization of polyphenol oxidase and photosystem II proteins. *Plant Physiology* **96**, 26–31.
- Lee B, Kim K, Jung W, Avice J, Ourry A, Kim T.** 2007. Peroxidases and lignifications in relation to the intensity of water-deficit stress in white clover (*Trifolium repens* L.). *Journal of Experimental Botany* **58**, 1271–1279.
- Lee MRF, Winters AL, Scollan ND, Dewhurst RJ, Theodorou MK, Minchin FR.** 2004. Plant-mediated lipolysis and proteolysis in red clover with different polyphenol oxidase activities. *Journal of the Science of Food and Agriculture* **84**, 1639–1645.
- Lee MRF, Tweed JKS, Minchin FR, Winters AL.** 2009. Red clover polyphenol oxidase: Activation, activity and efficacy under grazing. *Animal Feed Science and Technology* **149**, 250–264.
- Leufken CM, Moerschbacher BM, Dirks-Hofmeister ME.** 2014. Dandelion PPO-1/PPO-2 domain swaps: the C-terminal domain modulates the pH optimum and the linker affects SDS-mediated activation and stability. *Biochimica et Biophysica Acta* **1854**, 178–186.
- Li L, Steffens JC.** 2002. Overexpression of polyphenol oxidase in transgenic tomato plants results in enhanced bacterial disease resistance. *Planta* **215**, 239–247.
- Liu Y, Gao L, Xia T, Zhao L.** 2009. Investigation of the site-specific accumulation of catechins in the tea plant (*Camellia sinensis* (L.) O. Kuntze) via vanillin–HCl staining. *Journal of Agricultural and Food Chemistry* **57**, 10371–10376.
- Marusek CM, Trobaugh NM, Flurkey WH, Inlow JK.** 2006. Comparative analysis of polyphenol oxidase from plant and fungal species. *Journal of Inorganic Biochemistry* **100**, 108–123.
- Martinez VM, Whitaker JR.** 1995. The biochemistry and control of enzymatic browning. *Trends in Food Science and Technology* **6**, 195–200.
- Mayer AM.** 2006. Polyphenol oxidases in plants and fungi: going places? A review. *Phytochemistry* **67**, 2318–2331.
- Mayer AM, Harel E.** 1979. Polyphenol oxidases in plants. *Phytochemistry* **18**, 193–215.
- Meyer HU, Biehl B.** 1980. Activities and multiplicity of phenolase from spinach chloroplasts during leaf aging. *Phytochemistry* **19**, 2267–2272.
- Molitor C, Mauracher S, Kaintz C.** 2013. Crystal structures of the latent and *in vivo* proteolytically activated aurone synthase from *Coreopsis lanceolata*. In: *Konferenzband*. 15: Österreichische Chemietage, Graz, Austria, 23–26 September 2013.
- Nakayama T, Yonekura-Sakakibara K, Sato T, et al.** 2000. Aureusidin synthase: a polyphenol oxidase homolog responsible for flower coloration. *Science* **290**, 1163–1166.
- Nakayama T, Sato T, Fukui Y, Yonekura-Sakakibara K, Hayashi H, Tanaka Y, Kusumi T, Nishino T.** 2001. Specificity analysis and mechanism of aurone synthesis catalyzed by aureusidin synthase, a polyphenol oxidase homolog responsible for flower coloration. *FEBS Letters* **499**, 107–111.
- Neill SO, Gould KS.** 2003. Anthocyanines in leaves: light attenuators or antioxidants? *Functional Plant Biology* **30**, 865–873.
- Neumann J, Drechsler Z.** 1967. Inhibition of photo-induced electron transport and related reactions in isolated chloroplasts by phenol. *Plant Physiology* **42**, 573–577.
- O'Brien PJ.** 1991. Molecular mechanisms of quinone cytotoxicity. *Chemico-Biological Interactions* **80**, 1–41.
- Ono E, Hatayama M, Isono Y, et al.** 2006. Localization of a flavonoid biosynthetic polyphenol oxidase in vacuoles. *The Plant Journal* **45**, 133–143.
- Parveen I, Threadgill MD, Moorby JM, Winters A.** 2010. Oxidative phenols in forage crops containing polyphenol oxidase enzymes. *Journal of Agricultural and Food Chemistry* **58**, 1371–1382.
- Pietta P.** 2000. Flavonoids as antioxidants. *Journal of Natural Products* **63**, 1035–1042.
- Rivero RM, Ruiz JM, García PC, López-Lefebvre L, Sánchez E, Romero L.** 2001. Resistance to cold and heat stress: accumulation of phenolic compounds in tomato and watermelon plants. *Plant Science* **160**, 315–321.
- Rocha AMCN, Morais AMMB.** 2001. Characterization of polyphenoloxidase (PPO) extracted from 'Jonagored' apple. *Food Control* **12**, 85–90.
- Satō M.** 1966. Metabolism of phenolic substances by the chloroplasts. II. Conversion by the isolated chloroplasts of *p*-coumaric acid to caffeic acid. *Phytochemistry* **5**, 385–389.
- Saunders JA, McClure JW.** 1976a. The distribution of flavonoids in chloroplasts of twenty five species of vascular plants. *Phytochemistry* **15**, 809–810.
- Saunders JA, McClure JW.** 1976b. The occurrence of photoregulation of flavonoids in barley plastids. *Phytochemistry* **15**, 805–807.
- Schmitz GE, Sullivan ML, Hatfield RD.** 2008. Three polyphenol oxidases from red clover (*Trifolium pratense*) differ in enzymatic activities and activation properties. *Journal of Agricultural and Food Chemistry* **56**, 272–280.
- Sheptovitsky YG, Brudvig GW.** 1996. Isolation and characterization of spinach photosystem II membrane-associated catalase and polyphenol oxidase. *Biochemistry* **35**, 16255–16263.
- Sofo A, Dichio B, Xiloyannis C, Masia A.** 2005. Antioxidant defences in olive trees during drought stress: changes in activity of some antioxidant enzymes. *Functional Plant Biology* **32**, 45–53.
- Solomon EI, Sundaram UM, Machonkin TE.** 1996. Multicopper oxidases and oxygenases. *Chemical Reviews* **96**, 2563–2605.
- Sommer A, Ne'eman E, Steffens JC, Mayer AM, Harel E.** 1994. Import, targeting, and processing of a plant polyphenol oxidase. *Plant Physiology* **105**, 1301–1311.
- Steffens JC, Harel E, Hunt MD.** 1994. Polyphenol oxidase. In: Ellis BE, Kuroki GW, Stafford HA, eds. *Genetic engineering of plant secondary metabolism*. New York: Plenum Publishing Corporation, 275–312.
- Subramanian N, Venkatesh P, Ganguli S, Sinkar VP.** 1999. Role of polyphenol oxidase and peroxidase in the generation of black tea theaflavins. *Journal of Agricultural and Food Chemistry* **47**, 2571–2578.
- Sullivan ML, Hatfield RD, Thoma SL, Samac DA.** 2004. Cloning and characterization of red clover polyphenol oxidase cDNAs and expression of active protein in *Escherichia coli* and transgenic alfalfa. *Plant Physiology* **136**, 3234–3244.
- Sullivan ML, Hatfield RD, Samac DA.** 2008. Cloning of an alfalfa polyphenol oxidase gene and evaluation of its potential in preventing postharvest protein degradation. *Journal of the Science of Food and Chemistry* **88**, 1406–1414.
- Sullivan ML, Zarnowski R.** 2010. Red clover coumarate 3'-hydroxylase (CYP98A44) is capable of hydroxylating *p*-coumaroyl-shikimate but not *p*-coumaroyl-malate: implications for the biosynthesis of phaselic acid. *Planta* **231**, 319–328.
- Thipyapong P, Joel DM, Steffens JC.** 1997. Differential expression and turnover of the tomato polyphenol oxidase gene family during vegetative and reproductive development. *Plant Physiology* **113**, 707–718.

- Thipyapong P, Hunt MD, Steffens JC.** 2004a. Antisense downregulation of polyphenol oxidase results in enhanced disease susceptibility. *Planta* **220**, 105–117.
- Thipyapong P, Melkonian J, Wolfe DW, Steffens JC.** 2004b. Suppression of polyphenol oxidases increases stress tolerance in tomato. *Plant Science* **167**, 693–703.
- Thipyapong P, Stout MJ, Attajarusit J.** 2007. Functional analysis of polyphenol oxidases by antisense/sense technology. *Molecules* **12**, 1569–1595.
- Tolbert NE.** 1973. Activation of polyphenol oxidase of chloroplasts. *Plant Physiology* **51**, 234–244.
- Tóth SZ, Schansker G, Garab G.** 2013. The physiological roles and metabolism of ascorbate in chloroplasts. *Physiologia Plantarum* **105**, 1301–1311.
- Tran LT, Taylor JS, Constabel CP.** 2012. The polyphenol oxidase gene family in land plants: lineage-specific duplication and expansion. *BMC Genomics* **13**, 395.
- Trebst A, Eck H, Wagner S.** 1963. Effects of quinones and oxygen in the electron transport system of chloroplasts. In: National Research Council Committee on Photobiology, ed. *Photosynthetic mechanisms of green plants*. Washington, US: National Academy of Sciences, publication 1145, 174–194.
- Trebst A, Depka B.** 1995. Polyphenol oxidase and photosynthesis research. *Photosynthesis Research* **46**, 41–44.
- Vaughn KC, Duke SO.** 1984. Function of polyphenol oxidases in higher plants. *Physiologia Plantarum* **60**, 106–112.
- Veljovic-Jovanovic S, Kukavica B, Navari-Izzo F.** 2008. Characterisation of polyphenol oxidase changes induced by dessication of *Ramonda serbica* leaves. *Physiologia Plantarum* **132**, 407–416.
- Vinyard DJ, Ananyev GM, Dismukes GC.** 2013. Photosystem II: the reaction center of oxygenic photosynthesis. *Annual Review of Biochemistry* **82**, 577–606.
- Webb KJ, Cookson A, Allison G, Sullivan M, Winters AL.** 2013. Gene expression patterns, localization and substrates of polyphenol oxidase in red clover (*Trifolium pratense* L.). *Journal of Agricultural and Food Chemistry* **61**, 7421–7430.
- Webb KJ, Cookson A, Allison G, Sullivan M, Winters AL.** 2014. Polyphenol oxidase affects normal nodule development in red clover (*Trifolium pratense* L.). *Frontiers in Plant Science* doi: 10.3389/fpls.2014.00700
- Winters AL, Minchin FR, Merry RJ, Morris P.** 2003. Comparison of polyphenol oxidase activity in red clover and perennial ryegrass. In: Abberton MT, Andrews M, Skot L, Theodorou MK, eds. *Aspects 70: Crop quality: its role in sustainable livestock production*. Association of Applied Biologists Conference, Manchester, 15–16 December 2003, 121–128.
- Winters AL, Minchin FR, Michaelson-Yeates TPT, Lee MRF, Morris P.** 2008. Latent and active polyphenol oxidase (PPO) in red clover (*Trifolium pratense*) and use of a low PPO mutant to study the role of PPO in proteolysis reduction. *Journal of Agricultural and Food Chemistry* **56**, 2817–2824.
- Yoruk R, Marshall MR.** 2003. Physicochemical properties and function of plant polyphenol oxidase: a review. *Journal of Food Biochemistry* **27**, 361–422.